



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>A61K 7/32, 7/46</b>	<b>A1</b>	(11) International Publication Number: <b>WO 99/44575</b> (43) International Publication Date: 10 September 1999 (10.09.99)
<p>(21) International Application Number: PCT/US99/04147</p> <p>(22) International Filing Date: 25 February 1999 (25.02.99)</p> <p>(30) Priority Data: 60/076,861 5 March 1998 (05.03.98) US 09/213,625 18 December 1998 (18.12.98) US</p> <p>(71) Applicant: COLGATE-PALMOLIVE COMPANY [US/US]; 300 Park Avenue, New York, NY 10022 (US).</p> <p>(72) Inventors: GUENIN, Eric, P.; 305 Kentshire Court, Hopewell Township, NJ 08638 (US). BOUDOT, Pierre, Gabriel; 17, avenue des Champs Pierreux, F-92000 Nanterre (FR). SIL-LON, Pascal, Michel, Pierre; 17, avenue des Champs Pierreux, F-92000 Nanterre (FR). VINCENTI, Paul, Joseph; 95 Minnisink Road, Jefferson, NJ 07849 (US). TAYLOR, C., Donald; 10 Wheeler Road, Kendall Park, NJ 08824 (US). DURAND, Philippe, Michel; 136 North Road, Hythe, Kent CT21 4AT (GB).</p> <p>(74) Agent: MIANO, Rosemary, M.; Colgate-Palmolive Company, 909 River Road, P.O. Box 1343, Piscataway, NJ 08855-1343 (US).</p>		<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p><b>Published</b> <i>With international search report.</i></p>
<p>(54) Title: FRAGRANCE ENHANCING COMPOSITIONS FOR COSMETIC PRODUCTS</p> <p>(57) Abstract</p> <p>This invention relates to fragrance enhancing compositions which are capable of controlling malodor from a human body to a significant extent thereby reducing the overall amount of fragrance required to achieve a satisfactory cosmetic product, especially an underarm product. The fragrance enhancing compositions of this invention are made by combining at least three components from a selected group of non-nitromusk materials.</p> <p style="text-align: right;"><b>BEST AVAILABLE COPY</b></p>		

*FOR THE PURPOSES OF INFORMATION ONLY*

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

## FRAGRANCE ENHANCING COMPOSITIONS FOR COSMETIC PRODUCTS

Field of the Invention

This invention relates to fragrance enhancing compositions which may be used  
5 in a variety of products, especially in underarm products such as deodorants and  
antiperspirants, to give a deodorant effect against malodor from underarm perspiration  
and enhance the performance of the fragrance components of such products.

Background of the Invention

A variety of approaches have been and continue to be taken for reducing and  
10 or eliminating human body malodor, especially from underarm perspiration. These  
approaches include reducing/eliminating perspiration itself, reducing/eliminating odor  
at the root cause (for example, interfering with the degradation of perspiration on skin  
caused by the action of bacteria) as well as masking the odors themselves.

U.S. Patent 5,540,853 to Trinh et al describes personal treatment compositions  
15 and cosmetic compositions containing enduring perfume wherein the enduring  
perfume is evaluated by a "calculated log P" parameter. PCT application WO  
97/30689 to Trinh et al describes personal treatment compositions which can be used  
as leave-on products which are evaluated by the same criteria.

EP patent application 0 760 243 A1 to Groverman et al describes the use of  
20 allylic perfumes as a malodor reduction agent.

EP patent application 0 545 556 A2 to Behan et al teaches a perfume  
composition containing at least 50% by weight of specific components (in minimum  
percentages of each component) selected from the group consisting of ethers,

salicylates, alcohols, acetate/propionate esters and methyl ethyl ketones where members of at least 4 of the groups must be included.

U.S. Patent Number 5,601,809 teaches the neutralization of axillary malodor.

Various references which describe evaluation of underarm deodorants or  
5 methods of evaluating products used as deodorants include P.M. Baxter et al,  
International Journal of Cosmetic Science, Volume 5, 85-95 (1983) ; Chemical  
Senses, Volume 13, No. 3, 463-471 (1988); J.A. Killian, J. Soc. Cosmetic Chemists,  
Volume 3, 30-76 (1952), incorporated by reference in their entirety herein.

Specific studies of selected fragrance agents include A. Baydar et al, Chemical  
10 Senses, Volume 18, No. 6, 661-668 (1993) (olfactory threshold for androstenone and  
galaxolide); Chemical Senses, Volume 18, No. 3 245-256 (1993) (mutual cross  
adaptation of the volatile steroid androstenone and a non-steroid perceptual analog);  
L.J. Flett, J. Soc. Cosmetic Chemists, Volume 1, 304-310 (1949) deodorant properties  
of nacconol).

15 U.S. Patent 5,354,737 to Barr et al describes selected fragrancing  
compositions such as Tonalid in deodorant products.

There is still a need, however, for improving the way that cosmetic  
compositions are fragranced. Thus it is an object of the invention to provide  
fragrancing compositions which are useful in controlling malodor, especially  
20 underarm malodor. It is another object of this invention to provide fragrance  
enhancing components of a fragrance system for cosmetic products so that the amount  
of the overall fragrance components can be reduced. It is a further object of the  
invention to provide fragrance enhancing components that improves the performance

of conventional fragrancing products. It is still another object of the invention to reduce the overall irritation potential of cosmetic products by reducing the amount of fragrance needed to provide a satisfactory product. It is also an object of the invention to provide fragrancing compositions which exhibit improved ability to mask underarm odor. It is yet another object of the present invention to provide improved underarm products such as deodorants and antiperspirants containing such fragrancing compositions. These and other objects of the invention will become apparent from the following description of the invention.

#### Summary of the Invention

10        This invention relates to fragrance enhancing compositions which are capable of controlling malodor from a human body to a significant extent thereby reducing the overall amount of fragrance required to achieve a satisfactory cosmetic product, especially an underarm product. The fragrance enhancing compositions of the invention are formulated with selected components with high olfactory impact  
15        properties and good substantivity (for example, 24 hour endurance) which are combined for controlling human malodor, especially underarm sweat malodor. The fragrances made with the fragrance enhancing compositions of this invention are long lasting (enduring) as evaluated by accepted panel evaluations. By using the compositions of this invention it may also be possible to lower the amount of  
20        fragrance normally used in such cosmetic products (usually in the range of 0.1-10% by weight based on the total weight of the cosmetic composition) to some lesser amount.

The fragrance enhancing compositions of this invention are made by combining at least three components from the group consisting of the following non-nitromusk materials:

- 5 (1) Cinnamon Leaf Oil Ceylan {a Natural Essential Oil - predominantly 2-methoxy 4 allyl phenol};
- (2) Vetiver Oil Java {a Natural Essential Oil - predominantly Vetiverol alcohol};
- (3) Orange Oil Morocco {a Natural Essential Oil - predominantly d-Limonene};
- (4) Patchouli Oil {a Natural Essential Oil - predominantly Patchouli Alcohol and Patchoulene};
- 10 (5) Iso Methyl Cedryl Ketone A {a multi-component proprietary formulation available from Fragrance Resources, Keyport, New Jersey; {this material is predominantly comprised of esters with additional minor amounts of alcohols, essential oils, terpenes and others};
- (6) Pelargonyl {a multi-component proprietary formulation available from Synarome,
- 15 Bois Colombes, France; this material is comprised of mostly ethers with some portions of esters, alcohols and terpenes};
- (7) Cassis 345 B {a multi-component proprietary formulation available from Firmenich SA, Geneva, Switzerland; this material is primarily composed of alcohols and esters with minor amounts of natural extracts from essential oils};
- 20 (8) nonyl aldehyde;
- (9) decyl aldehyde;

- (10) Ambroxan® {8- $\alpha$ , 12-oxido-13,14,15,16-tetranorlabdane or 3aR-(3a, $\alpha$ ,5a, $\alpha$ ,9a, $\beta$ ,9b, $\beta$ ))-dodecahydro-3a,6,6,9a-tetramethylnaphtho(2,1-b)furan};
- (11) Alpha-Iso-Amyl-Cinnamic {amyl cinnamic aldehyde};
- 5 (12) benzyl propionate;
- (13) Carvone Laevo {L-1-methyl-4-isopropenyl-6-cyclohexene-2-one};
- (14) Beta-Gamma Hexenyl Salicylate {cis-3-hexenyl salicylate};
- (15) Citral {3,7-dimethyl-2,6-octadienal};
- (16) Citronellol Dextro {3-7-dimethyl-6-octen-1-ol};
- 10 (17) Cyclamen Aldehyde {2-methyl-3-(para-isopropylphenyl) propionaldehyde};
- (18) Damascone Alpha {2-buten-1-one, 1-(2,6,6-trimethyl-2-cyclohexen-1-yl)};
- (19) Decyl Aldehyde Natural {Decanal Natural};
- (20) Ethyl Vanillin {3-ethoxy-4-hydrobenzaldehyde};
- (21) Eugenol {2-methoxy-4-allyl phenol};
- 15 (22) Florazon {4-ethyl- $\alpha$ ,  $\alpha$ -dimethyl-benzenepropanal};
- (23) Geraniol {2-trans-3,7-dimethyl-2,6-octadien-8-ol};
- (24) Herbanate {3-(1-methyl-ethyl) bicyclo (2,2,1) hept-5-ene-2-carboxylic acid ethyl ester};
- (25) Iso Butyl Quinoline {6-secondary butyl quinoline};
- 20 (26) Menthol Laevo {5-methyl-2-isopropyl cyclohexanol};
- (27) Menthone Laevo {4-isopropyl-1-methyl cyclohexan-3-one};
- (28) Iralia Total® {Methyl Ionone (mixture of isomers)};

- (29) Methyl Ionone Gamma Coeur {3-buten-2-one, 3 methyl-4-(2,6,6-trimethyl-2-cyclohex-1-yl)};
- (30) para-hydroxy phenyl butanone crystals {4-(4-hydroxyphenyl)-2-butanone};
- (31) 4-methyl-2-phenyl-3,6-dihydropyran;
- 5 (32) Rootanol 100® {2 -(1,1-dimethylethyl)-4-methyl-cyclohexanol};
- (33) Terpineol Alpha {1-methyl-4-isopropyl-1-cyclohexen-8-ol};
- (34) Vanillin {4-hydroxy-3-methoxy benzaldehyde}; and
- (35) 2,4-diethoxy-5-methyl- pyrimidine}.

#### Detailed Description of the Invention

10 The term "Deo-Key™" is applied here to fragrance enhancing compositions which provide an improved way to control malodor, especially underarm malodor either alone or in combination with other fragrances of the type conventionally used in cosmetic compositions, especially those used in underarm products. The fragrance enhancing compositions (also called fragrance enhancing components herein) are

15 made by combining at least three (and more particularly five) members of the group consisting of the thirty-five ingredients listed above in proportions as follows wherein the amounts are based on the total amount of the fragrance component in the cosmetic product:

- (a) 0.5-1.0% Cinnamon Leaf Oil Ceylan;
- 20 (b) 0.01-0.3% Vetiver Oil Java;
- (c) 0.5-6.0% Orange Oil Morocco;
- (d) 0.05-6.0% Patchouli Oil;



- (e) 0.05-4.0% Iso Methyl Cedryl Ketone A;
- (f) 0.05-6.0% Pelargonyl;
- (g) 0.05-0.9% Cassis 345 B;
- (h) 0.005-0.05% nonyl Aldehyde;
- 5 (i) 0.005-0.05% decyl Aldehyde;
- (j) 0.05-0.30% 3aR-(3a, alpha, 5a, alpha, 9a, beta, 9b, beta))-dodecahydro-3a, 6, 6, 9a-tetramethyl-naphthto(2, 1-b)furan;
- (k) 0.05-10.0% amyl cinnamic aldehyde;
- (l) 0.05-0.9% benzyl propionate;
- 10 (m) 0.05-0.30% L-1-methyl-4-isopropenyl-6-cyclohexene-2-one;
- (n) 0.05-0.30% cis-3-hexenyl salicylate;
- (o) 0.05-0.70% 3,7-dimethyl-2,6-octadienal;
- (p) 0.05-2.00% 3-7-dimethyl-6-octen-1-ol;
- (q) 0.05-0.9% 2-methyl-3-(para-isopropylphenyl) propionaldehyde;
- 15 (r) 0.05-0.9% 2-buten-1-one, 1-(2,6,6-trimethyl-2-cyclohexen-1-yl);
- (s) 0.005-0.05% decyl aldehyde Natural;
- (t) 0.05-0.3% 3-ethoxy-4-hydrobenzaldehyde;
- (u) 0.05-1.00% 2-methoxy-4-allyl phenol;
- (v) 0.05-0.30% 4-ethyl-alpha, alpha-dimethyl- benzenepropanal;
- 20 (w) 0.05-6.0% 2-trans-3,7-dimethyl-2,6-octadien-8-ol;
- (x) 0.05-4.0% 3-(1-methyl-ethyl) bicyclo (2,2,1) hept-5-ene-2-carboxylic acid ethyl ester;

- (y) 0.05-0.30% isobutyl quinoleine;
- (z) 0.05-2.0% 5-methyl-2-isopropyl cyclohexanol;
- (aa) 0.05-0.3% 4-isopropyl-1-methyl cyclohexan-3-one;
- (bb) 0.05-2.0% methyl ionone (mixture of isomers);
- 5 (cc) 0.05-2.0% 3-buten-2-one, 3 methyl-4-(2,6,6-trimethyl-2-cyclohex-1-yl;
- (dd) 0.05-0.90% para-hydroxy phenyl butanone crystals;
- (ee) 0.05-4.0% 4-methyl-2-phenyl-3,6-dihdropyrane;
- (ff) 0.05-0.30% 2-(1,1-dimethylethyl)-4-methyl-cyclohexanol;
- (gg) 0.05-6.0% 1-methyl-4-isopropyl-1-cyclohexen-8-ol;
- 10 (hh) 0.05-0.3% 4-hydroxy-3-methoxy benzaldehyde; and
- (ii) 0.05-0.3% 2,4-diethoxy-5-methyl-pyrimidine.

For cosmetic compositions an amount of 1.8-100%, particularly 1.8-60%, more particularly 1.8- 35%, with a specific range being 1.8-32.5%, and more specific ranges being 2.5-30%, 25-27%, and 20% by weight based on the total amount of

15 fragrance component added to make the cosmetic composition can be used to achieve the desired effect. It is also possible that a major portion of the fragrance portion such as 60-90% of the fragrance portion and even 100% of the fragrance portion is made of the fragrance enhancing compositions described for this invention.

The use of the fragrance enhancing compositions make the use of conventional

20 fragrances more effective in controlling underarm malodor and may also result in the ability to use lower amounts of fragrances to achieve performance equivalent to compositions not containing such fragrance-enhancing compositions. Although it is possible to use the fragrance enhancing compositions of this invention as the sole

fragrance in cosmetic products such as deodorants and/or antiperspirants, it has been found preferable (for aesthetic reasons to give a more rounded fragrance) to use at least a portion of conventional fragrances in the cosmetic compositions.

The conventional fragrances that can be combined with the fragrance enhancing compositions of the invention include those described in PCT application WO 97/30689 and incorporated by reference in its entirety herein. Specific non-limiting examples of suitable fragrances include those selected from the group consisting of those described therein. Specific groups include:

(a) esters of salicylic acid such as hexyl salicylate, hexenyl salicylate, isoamyl salicylate, benzyl salicylate and cyclohexyl salicylate;

(b) esters of cinnamic acid such as amyl cinnamate, cinnamyl cinnamate and methyl cinnamate;

(c) miscellaneous esters such as allyl cyclohexane propionate, amyl benzoate, para-tertiarybutylcyclohexyl acetate, cedryl acetate, cedryl formate, dihydro-isojasmonate, ethylene brassylate, ethyl undecylenate, geranyl anthranilate, geranyl phenyl acetate, linalyl benzoate, benzyl acetate, linalyl acetate, vetiveryl acetate;

(d) aldehydes such as amyl cinnamic aldehyde, cyclamen aldehyde, lillial, benzaldehyde, citronnellal, hydroxycitronellal;

(e) alcohols such as geraniol, linalool, nerol, phenyl ethyl alcohol, alpha-terpineol, eugenol, isoeugenol, alpha-citronellol, dihydromyrcenol, auranliol, cedrol, phenyl heptanol, phenol hexanol, alpha-santalol, undecavertol (4-methyl-3-decen-5-ol), benzyl alcohol;

(f) ketones such as benzophenone, dodecalactone, gamma-n-methyl ionone, delta-undecalactone, gamma-undecalactone, laevo-carvone, beta-methylnaphthyl ketone;

(g) nitromusk such as musk ketone, musk tibetine, musk indanone.

5       The individual components listed above used to form the fragrance enhancing compositions of this invention are commercially available and may be obtained from suppliers of fragrances and specialty chemicals who are known to those skilled in the art.

      The ability of the fragrance enhancing components of the invention to improve  
10   the performance of conventional fragrances may in some cases be due to synergy between the various components of the overall fragrance component (the total of all fragrance ingredients used in a cosmetic product).

      Each of the thirty-five components listed above have a sensory sniff rating as evaluated by panel tests. Such tests are described in the art and are well accepted as  
15   being appropriate ways of evaluating overall fragrance impact and effect. An example of one such test is described in International Journal of Cosmetic Science, Volume 5, pages 85-95 (1983). In such tests numerical rating scales are used to establish the relative performance of each fragrance. One such scale is, for example 0-5 (0 means a high performance, 5 a poor performance). On such a scale the individual  
20   compositions listed above as the group from which the fragrance enhancing compositions would be made would each have a rating between 0-2. Such fragrance raw materials improve deodorancy and longevity of cosmetic products (for example, deodorant sticks) in conjunction with fragrance aesthetics. Since fragrances are

sometimes responsible for skin irritation, it is also a benefit of the present invention that the overall amount of fragrance in a cosmetic products can be reduced by the use of the compositions of this invention.

The cosmetic compositions can include one or more active ingredients selected from, for example, a member selected from the group consisting of antiperspirant active materials, deodorant active materials, sunscreens, insect repellents, antifungal agents, antimicrobials (also called bacteriostats or antibacterials), and additional fragrances.

Where the composition contains an antiperspirant active, any of the known antiperspirant active materials can be utilized. These can be added as solutions, suspensions or directly during mixing. These include, by way of example (and not of a limiting nature), aluminum chlorohydrate, aluminum chloride, aluminum sesquichlorohydrate, zirconyl hydroxychloride, aluminum-zirconium glycine complex (for example, aluminum zirconium trichlorohydrate gly, aluminum zirconium pentachlorohydrate gly, aluminum zirconium tetrachlorohydrate gly and aluminum zirconium octochlorohydrate gly), aluminum chlorohydrate PG, aluminum chlorohydrate PEG, aluminum dichlorohydrate PG, and aluminum dichlorohydrate PEG. The aluminum-containing materials can be commonly referred to as antiperspirant active aluminum salts. Generally, the foregoing metal antiperspirant active materials are antiperspirant active metal salts. In the embodiments which are antiperspirant compositions according to the present invention, such compositions need not include aluminum-containing metal salts, and can include other antiperspirant active materials, including other antiperspirant active metal salts.

Generally, Category I active antiperspirant ingredients listed in the Food and Drug Administration's Monograph on antiperspirant drugs for over-the-counter human use can be used. In addition, any new drug, not listed in the Monograph, such as aluminum nitratohydrate and its combination with zirconyl hydroxychlorides and nitrides, or aluminum-stannous chlorohydrates, can be incorporated as an antiperspirant active ingredient in antiperspirant compositions according to the present invention.

Antiperspirant actives can be incorporated into compositions according to the present invention in amounts in the range of 0.1 - 30%, preferably 15 - 25%, by weight, of the total weight of the composition. The amount used will depend on the formulation of the composition. For example, at amounts in the lower end of the broader range (for example, 0.1 - 10%), the antiperspirant active material will not substantially reduce the flow of perspiration, but will reduce malodor, for example, by acting as an antimicrobial material.

For embodiments of the invention which contain an antiperspirant (either at a level denominated "deodorant" or at a level denominated "antiperspirant") it is preferred that a stabilizing agent also be included. Examples of suitable stabilizing agents include cosmetically acceptable alkali metal salts, bases, amines and other nitrogen containing compounds, particularly guanidine carbonate (described in U.S. Patent 5,490,979 and assigned to the same owner as this application).

When a bacteriostat composition is included in the cosmetic products of this invention such known bacteriostats may include bacteriostatic quaternary ammonium compounds (such as cetyl-trimethylammonium bromide), 2-amino-2-methyl-1-

propanol (AMP), cetyl pyridinium chloride, 2, 4, 4'-trichloro-2'-hydroxydiphenylether (Triclosan), N-(4-chlorophenyl)-N'-(3,4-dichlorophenyl)urea (Triclocarban) and various zinc salts (for example, zinc ricinoleate). The bacteriostat can, illustratively, be included in the composition in an amount of 0.01-1.0% by weight, of the total weight of the composition. Triclosan, can illustratively be included in an amount of from 0.05% to about 0.5% by weight, of the total weight of the composition.

If sunscreens are used conventional agents may be included.

The base for cosmetic compositions made with the fragancing materials of this invention include soap based deodorants, especially those made with one or more glycols (propylene glycol, dipropylene glycol, tripropylene glycol, tetrapropylene glycol and/or other polypropylene glycols); silicone based deodorants and antiperspirants, especially those made with fatty alcohols.

Products made as cosmetic compositions may include antiperspirants and/or deodorants, deodorant soaps, body washes, creams, lotions, scented cosmetic products such as colognes and perfumes.

In one embodiment for a deodorant stick with such efficacy, fragrance raw materials should be used in an amount between 0.001% and 25.00% of the fragrance (neat composition) portion of the composition, particularly between 0.001% and 20.00%, more preferably between 0.05% and 15.00% of the fragrance, and even more preferably between 0.50% and 10.00% of the fragrance.

Deo-Key <sup>TM</sup> compositions combine fragrance raw materials such as examples described below. The examples listed below describe compositions used in a range of

- 1.8-32.6% based on the total fragrance component. For example 30% of a fragrance used at a 10% level would give a maximum level for the Deo- Key composition of 3.00% by weight of the overall cosmetic product. The following embodiments show how ranges of various fragrance components can be combined wherein the amounts listed are percents of the total fragrance component (unless another standard is listed such as "overall product").

	<u>Green Deo-Key™ Fragrance Enhancer</u>	<u>Range A</u>	<u>Range B</u>
	Cassis 345 B	0.20-1.50	0.40-0.90
	Damascone Alpha	0.20-1.50	0.40-0.90
10	Para Hydroxy Phenyl Butanone Crystals	0.20-1.50	0.40-0.90
	Herbanate	1.00-5.00	2.00-4.00
	Rosyrane	1.00-5.00	2.00-4.00
	Total	2.60-14.50	5.2-10.7

- 15 In another embodiment the Green Deo-Key fragrance can be used in an amount of from 25-27% of the total fragrance component (which is greater than the specifically described 2.60-14.50 amount), provided that the ratios of the individual ingredients are maintained in the same ranges as described for the 2.60-14.50 composition).

	<u>Wood Deo-Key™ Fragrance Enhancer</u>	<u>Range A</u>	<u>Range B</u>
20	Vanillin	0.02-0.50	0.05-0.30
	Vethymine	0.02-0.50	0.05-0.30
	Ambroxan	0.02-0.50	0.05-0.30
25	Benzenepropanal, 4 ethyl-alpha, alpha-dimethyl {Florazon 3/034395}	0.02-0.50	0.05-0.30



	Vetiver Oil Java	0.02-0.50	0.05-0.30
	Iso Methyl Cedryl Ketone A	1.00-5.00	2.00-4.00
	Terpineol Alpha	1.00-5.00	2.00-4.00
	Patchouli Oil	2.00-8.00	3.00-6.00
5	Amyl Cinnamic Aldehyde	4.00-12.00	6.00-10.00
	Total	8.10-32.50	13.2-25.5
	<u>Floral Wood Deo-Key™ Fragrance Enhancer</u>	<u>Range A</u>	<u>Range B</u>
	Damascone Alpha	0.02-0.50	0.05-0.30
	Rootanol 100	0.02-0.50	0.05-0.30
10	Carvone Laevo	0.02-0.50	0.05-0.30
	Benzyl Propionate	0.20-1.20	0.40-0.90
	Cyclamen Aldehyde	0.50-2.50	1.00-2.00
	Orange Oil Morocco	2.00-8.00	3.00-6.00
	Citronellol Dextro	2.00-8.00	3.00-6.00
15	Pelargonyl	2.00-8.00	3.00-6.00
	Total	6.76-29.20	10.6-21.8
	<u>Citrus Wood Deo-Key™ Fragrance Enhancer</u>	<u>Range A</u>	<u>Range B</u>
	Nonyl Aldehyde	0.002-0.1	0.005-0.05
	Decyl Aldehyde	0.002-0.1	0.005-0.05
20	Cis-3-Hexenyl Salicylate	0.02-0.4	0.05-0.30
	Citral Lemarome	0.02-0.4	0.05-0.30
	Cyclamen Aldehyde	0.20-1.20	0.40-0.90

	Citronellol Dextro	0.8-2.40	1.00-2.00
	Iralia Total	0.8-2.40	1.00-2.00
	Total	1.84-7.00	2.5-5.6
	<u>Amber Wood Deo-Key™ Fragrance Enhancer</u>	<u>Range A</u>	<u>Range B</u>
5	Isobutyl Quinoleine	0.02-0.4	0.05-0.30
	Ambroxan	0.02-0.4	0.05-0.30
	Cinnamon Leaf Oil Ceylan	0.4-1.20	0.60-1.00
	Eugenol	0.4-1.20	0.60-1.00
	Iso Methyl Cedryl Ketone A	1.00-5.00	2.00-4.00
10	Patchouli Oil	2.00-7.00	3.00-6.00
	Total	3.84-15.2	6.3-12.6
	<u>Mint Deo-Key™ Fragrance Enhancer</u>	<u>Range A</u>	<u>Range B</u>
	4-isopropyl-1-methyl cyclohexan-3-one {Menthone Laevo}	0.02-0.40	0.05-0.30
15	Ethyl Vanillin	0.02-0.40	0.05-0.30
	Decyl Aldehyde Natural {Decanal Natural 951512}	0.1-0.5	0.15-0.40
	5-Methyl 2-Isopropyl Cyclohexanol {Menthol Laevo}	0.8-2.40	1.00-2.00
	Methyl Ionone Gamma Coeur	0.8-2.40	1.00-2.00
20	Geraniol	2.00-7.00	3.00-6.00
	Total	3.74-13.10	5.3-11.0

It should be noted that other embodiments can contain more than the described totals for the compositions listed above, provided that the same ratios of the ingredients for each composition are maintained and a multiple of these ratios is used to give a total

amount greater than the specific embodiments listed above. This is particularly described above in the case of the Green Deo-Key composition.

The cosmetic compositions in which this invention can be used can take various forms including sticks, gels, soft solids, creams, liquids (for example, roll-  
5 ons), and aerosols.

### **SOLIDS**

	INGREDIENT	RANGES	PREFERRED RANGES
10	<b><u>Solid Stick "A"</u></b>		
	aluminum zirconium		
	tetrachlorohydrate-gly	15-25	18-22
	volatile silicone	30-60	45-55
15	(for example cyclomethicone)		
	stearyl alcohol	5-15	8-12
	talc	5-15	8-12
	hydrogenated castor oil	0-5	1-3
20	PPG-14 butyl ether	0-3	1-2
	fragrance component of which	0.1-10	1-3
	2.5- 30% is the Deo-Key™		
	composition (for example 2%		
	of the overall stick)		
25	alantoin	0.01-1	0.05-0.5
	glyceryl stearate	0.01-1	0.05-0.5
	PEG 100 stearate	0.01-1	0.05-0.5
	color (optional)	0.001-0.1	0.005-0.05
30	<b><u>Solid Stick "B"</u></b> (deodorant)		
	SD alcohol 40	10-80	40-70
	propylene glycol	5-20	10-15

	sodium stearate	2-10	6-8
	water	0-20	10-15
5	fragrance component of which 2.5- 30% is the Deo-Key™ composition	0.1-10	1-3
	trisodium EDTA	0.05-0.5	0.1-0.4
	color (optional)	0.001-0.01	0.005-0.01
	<u>Solid Stick C</u> (deodorant)		
10	Solvent (selected from a range of glycols as described for Stick "D")	5-88	60-75
	water	1-50	10-20
15	compatible gelling agent	1- 10	4-8
	emollient	0-5	1-2
	antibacterial agent	0.01-2.0	0.05-0.5
20	anti-irritancy agent	0.1-10	1-3
25	fragrance component of which 2.5- 30% is the Deo-Key™ composition (for example 2% of the overall stick)	0.1-10	0.5-3
	<u>Solid Stick D</u> (deodorant)		
30	glycol component (one or more of glycols such as propylene glycol, di-, tri-, tetra, and higher propylene glycols)	60-98	65-75
35	sodium stearate	0-10	6-8
	water	0-40	10-25
40	fragrance component of which 2.5- 30% is the Deo-Key™ composition (for example 2% of the overall stick)	0.1-10	0.5-3

GELS

	INGREDIENT	RANGES	PREFERRED RANGES
5	<u>Gel "A"</u>		
	cyclomethicone and dimethicone copolyol	5-30	8-10
	cyclomethicone	60-90	70-80
	aluminum chlorohydrate	10-25	
10	propylene glycol	10-20	12-18
	water	10-30	15-25
	fragrance component of which 2.5- 30% is the Deo-Key™ composition (for example 2% of the overall product)	0.5-10	1-3
15			

CREAMSCream "A"

	aluminum chlorohydrate	10-25	15-22
	aluminum chloride	5-15	6-12
20	water	40-70	50-65
	glyceryl stearate	1-20	5-15
	PEG-40 stearate	1-20	5-15
	cetyl palmitate or synthetic spermaceti	1-10	3-7
	glycerin	1-10	2-8
25	dimethicone	1-10	2-8
	isopropyl palmitate	1-10	2-5
	petrolatum and lanolin alcohol	1-10	2-6
	lanolin wax	1-10	2-8
	hydroxypropyl methylcellulose	0.5-1.5	0.8-1.2
30			

	fragrance component of which 2.5-30% is the Deo-Key™ composition (for example 1% of the overall product)	0.1-10	1-3
5	titanium dioxide	0.1-1	0.2-0.8
	lanolin	0.5-1	0.6-0.8
	propyl paraben	0.05-	0.1-0.2
	mineral oil	0.5-1	0.6-1
	<b><u>LIQUIDS</u></b>		
10	<b><u>Roll-on "A"</u></b> (antiperspirant)		
	aluminum chlorohydrate (for example 21%)	10-21	18-22
	water (for example 67%)	50-80	65-75
	PPG-11 stearyl ether (for example 7%)	5-10	6-8
	Steareth-2 (for example 5%)	2-10	4-8
15	Steareth-20 (for example 5%)	2-10	4-8
	fragrance component of which 2.5-30% is the Deo-Key™ composition (for example 1% of the overall product)	0.1-10	1-3
20	<b><u>Roll-on "B"</u></b> (antiperspirant)		
	zirconium-aluminum glycine hydroxychloride complex	10-25 (for example 20%)	18-22
	water	20-40 (for example 24.3%)	24-30
25	PEG-40 stearate	5-15 (for example 10%)	8-12
	glyceryl stearate	5-15 (for example 10%)	8-12
30	glycerine	5-10 (for example 6.8%)	6-8
	refined paraffin	4-10 (for example 6.2%)	5-9

	isopropyl palmitate	5-10 (for example 7.3%)	6-9
5	magnesium aluminum silicate	10-20 (for example 15.2%)	12-18
	fragrance component of which 2.5-30% is the Deo-Key™ composition (for example 1.5% of the overall stick)	0.1-10	1-3
10	<b><u>Roll-on "C"</u></b>		
	aluminum chlorohydrate	10-25 (for example 21%)	15-22
	cyclomethicone	40-80 (for example 70%)	60-70
15	quaternium-18 hectorite	0-6 (for example 4%)	3-5
	SD alcohol 40 (3.1% by volume)	1-4 (for example 2%)	2.5-3.5
20	Steareth-20	0-4 (for example 2%)	1-3
	fragrance component of which 2.5-30% is the Deo-Key™ composition (for example 1% of the overall product)	0.5-10	1-3
25	<b><u>Roll-on "D"</u></b>		
	INGREDIENT	RANGES	PREFERRED RANGES
	cyclomethicone	10-80	35-55
	cyclomethicone and dimethicone copolyol	5-20	10-15
30	aluminum zirconium tetrachlorohydrex-gly	10-25	15-22
	propylene glycol	5-15	7-12
	water	0-5	2-4
35			

fragrance component of  
which 2.5-30% is the  
Deo-Key™ composition

0.5-10

1-3

### AEROSOLS

#### 5 Aerosol "A"

An aerosol formulation made with the following amounts of ingredients based on the total weight of the composition: aluminum chlorohydrate (for example 10-22%); isobutane (for example 30-45%); cyclomethicone (for example 10-20%); isopropyl myristate (for example 2-5%); dimethicone (for example 2-4%); quaternium-18 hectorite (for example 1-2%); propylene carbonate (for example 0.5-1%); effective amount of a fragrance component of which 2.5-30% is the Deo-Key™ composition (for example 1% of the overall product).

#### Aerosol "B"

An aerosol formulation made with the following amounts of ingredients based on the total weight of the composition: SD alcohol 40 (for example 10-30%); isobutane (for example 10-30%); propane (for example 10-20%); propylene glycol (for example 5-20%); butane (for example 5-10%); effective amount of a fragrance component of which 2.5-30% is the Deo-Key™ composition (for example 1% of the overall product); effective amount of Triclosan.

#### 20 Pump Spray "A"

A pump spray formulation made with the following amounts of ingredients based on the total weight of the composition: cyclomethicone (for example 40-60%); SD alcohol 40B (anhydrous alcohol) (for example 20-40%); PEG-3 myristyl ether (for example 2-10%); C12-15 alcohol benzoates (for example 2-10%); water (for example 2-10%); effective amount of a fragrance component of which 2.5-30% is the Deo-Key™ composition; dimethicone (for example 0.5-5%); zinc phenosulfonate (for example 0.1-1.0%).



Pump Spray "B" - wherein amounts are based on the total weight of the composition.

5	INGREDIENT	RANGES OF INGREDIENTS	PARTICULAR EXAMPLE
	cyclomethicone	30-50%	33%
	mineral oil	1-10%	2.0%
	phenyltrimethicone	1-10%	5.0%
10	cyclomethicone and dimethicone copolyol	0.5-10%	10%
	polysorbate 20	0.5-5%	1.0%
	aluminum chlorohydrate	5-25%	10-25%
15	fragrance component of which 2.5-30% of the composition is the Deo-Key™ composition	0.1-10	

#### EXAMPLES

The following Examples are offered as illustrative of the invention and are not  
 20 to be construed as limitations thereon. In the Examples and elsewhere in the  
 description of the invention, chemical symbols and terminology have their usual and  
 customary meanings. Temperatures are in degrees C unless otherwise indicated. The  
 amounts of the components are in weight percents based on the standard described; if  
 no other standard is described then the total weight of the compositions is to be  
 25 inferred. Various names of chemical components include those listed in the CTFA  
International Cosmetic Ingredient Dictionary (Cosmetics, Toiletry and Fragrance  
 Association, Inc., 4<sup>th</sup> ed. 1991). Note that the antiperspirant actives described in the  
 examples and elsewhere are usually added in the form of a solution for example as a  
 50% solution. Effective amounts of fragrance component are in the range of 0.1-10%

by weight based on the total weight of the cosmetic composition, a portion of which is described as being the Deo-Key composition.

Example 1 - General Method A

The Deo-Key fragrance enhancing compositions of this invention may be made by conventional mixing techniques known to those skilled in the art. Such techniques will take into account the solid or liquid nature of the products involved and the need to pre-dissolve or melt ingredients before they are blended. The solid ingredients should be added first and dissolved or melted until a liquid is obtained by heating in a sand bath (for example, at a temperature in the range of 40 -60 degrees C). The liquid ingredients may then be added in a subsequent step and the final mixture will be completed at room temperature with stirring (for example, at 500 RPM with a stirring machine AM 3000 D from HEIDOLPH).

For example, in one method the ingredients are weighed in a beaker which has been tared (such as with a PJ3600 Delta Range balance from METTLER). The total quantity of Deo-Key material should be calculated to have a minimum quantity of individual material greater than 1 gram (minimum weighable quantity). If any error occurs it should be corrected by restarting the entire weighing process; thus, this sequential methodology of weighing in the same beaker avoids the loss of ingredients during transfer from one beaker to another.

20

Example 2

Test samples can be prepared by taking ten 6 cm diameter (2.5 inches) circles of Webril® pads and treating them with a selected amount of test product. The test sample is rubbed or applied on one side of the pad. Each pad is then placed in a

plastic disposable Petri dish which has been labeled. Each pad is allowed to dry for 24 hours at room temperature with the lid of the Petri dish removed. After the 24 hour period, a solution of synthetic malodor material is applied to each pad in an amount of about 9X excess of the sample test product by applying it to the treated surface of the pad. The solution is added dropwise to cover the entire surface of the treated pad. The sample is allowed to dry for an additional 2 hours before placing the lids on the Petri dishes. The plates are stored at room temperature until the 48 hour odor evaluation is performed. The odor assessment is done by a number of trained personnel, using some preset numerical scale.

10

Example 3Testing Method for Fragrance Effectiveness Against Malodor

The Green Deo-Key™ and Woody Deo-Key™ compositions prepared by the method described in Example 1 were tested against a traditional fragrance formulation of 2.5% made as follows. Evaluations of the effectiveness of each test sample were done by a panel using a rating system of 0-7 to evaluate the effectiveness of the sample in reducing malodor. The data shown below demonstrates that 0.4% levels of the Deo-Key™ compositions listed were as effective as traditional fragrance at a level of 2.5%. This indicates that the Deo-Key™ compositions may be used at much lower levels.

20

Type and Level of Deo-Key	Score	Control
0.4% green Deo-key	0.3	0.4
0.4% Woody Deo-key	0.6	0.6

Example 4

A comparative evaluation using a panel of five trained panelists was performed. The three samples tested were Sample A1 = Commercial Product A believed to contain about 2.5% of a conventional fragrance; Sample B1 = a cosmetic stick formulated with 0.4% of a Green Deo-Key composition as described above; and Sample C1 = the same cosmetic stick formulation as in Sample B but without the Deo-Key composition of the invention, but with 1.1% of a conventional fragrance. The evaluation results are given as an average of the ratings by the panel for perception of malodor. The higher the number the poorer the performance of the product: Sample A1 - 0.45; Sample B1 - 0.33; Sample C1 - 0.58.

Example 5

The procedure of Example 4 was repeated where Sample A2 = Sample A1; Sample B2 = Sample B1; and Sample C2 = the same cosmetic stick formulation as in B1 except that 0.6% of the Deo-Key composition was used. The evaluation data is as follows: Sample A2 - 0.45; Sample B2 - 0.33; Sample C2 - 0.19.

## We Claim

1. A fragrance enhancing component for a cosmetic composition made by adding as part of a total fragrance component at least three members selected from the group consisting of the following non-nitromusk materials:
  - 5 (a) Cinnamon Leaf Oil Ceylan;
  - (b) Vetiver Oil Java;
  - (c) Orange Oil Morocco;
  - (d) Patchouli Oil;
  - (e) Iso Methyl Cedryl Ketone A;
  - 10 (f) Pelargonyl;
  - (g) Cassis 345 B;
  - (h) Nonyl Aldehyde;
  - (i) Decyl Aldehyde;
  - (j) 3aR-(3a,α,5a,α,9a,β,9b,β))-dodecahydro-3a,6,6,9a-tetramethyl-
  - 15 naphthto(2,1-b)furan;
  - (k) amyl cinnamic aldehyde;
  - (l) benzyl propionate;
  - (m) L-1-methyl-4-isopropenyl-6-cyclohexene-2-one;
  - (n) cis-3-hexenyl salicylate;
  - 20 (o) 3,7-dimethyl-2,6-octadienal;
  - (p) 3-7-dimethyl-6-octen-1-ol;
  - (q) 2-methyl-3-(para-isopropylphenyl) propionaldehyde;

- (r) 1-(2,6,6-trimethyl-2-cyclohexen-1-yl-2-buten-1-one);
- (s) decyl aldehyde natural;
- (t) 3-ethoxy-4-hydrobenzaldehyde;
- (u) 2-methoxy-4-allyl phenol;
- 5 (v) benzenepropanal, 4-ethyl-alpha, alpha-dimethyl;
- (w) 2-trans-3,7-dimethyl-2,6-octadien-8-ol;
- (x) 3-(1-methyl-ethyl) bicyclo (2,2,1) hept-5-ene-2-carboxylic acid ethyl ester;
- (y) isobutyl quinoline;
- (z) 5-methyl-2-isopropyl cyclohexanol;
- 10 (aa) 4-isopropyl-1-methyl cyclohexan-3-one;
- (bb) methyl ionone (mixture of isomers);
- (cc) 3 methyl-4-(2,6,6-trimethyl-2-cyclohex-1-yl-3-buten-2-one);
- (dd) para-hydroxy phenyl butanone crystals;
- (ee) 4-methyl-2-phenyl-3,6-dihydropyran;
- 15 (ff) 2 -(1,1-dimethylethyl)-4-methyl-cyclohexanol;
- (gg) 1-methyl-4-isopropyl-1-cyclohexen-8-ol;
- (hh) 4-hydroxy-3-methoxy benzaldehyde; and
- (ii) 2,4-diethoxy-5-methyl-pyrimidine.

- 20           2.       A fragrance enhancing component according to Claim 1 made by combining at least three members from the group consisting of the following non-nitromusk materials:

- (a) 0.5-1.0% Cinnamon Leaf Oil Ceylan;
- (b) 0.01-0.3% Vetiver Oil Java;
- (c) 0.5-6.0% Orange Oil Morocco;
- (d) 0.05-6.0% Patchouli Oil;
- 5 (e) 0.05-4.0% Iso Methyl Cedryl Ketone A;
- (f) 0.05-6.0% Pelargonyl;
- (g) 0.05-0.9% Cassis 345 B;
- (h) 0.005-0.05% nonyl aldehyde;
- (i) 0.005-0.05% decyl aldehyde;
- 10 (j) 0.05-0.30% 3aR-(3a,α,5a,α,9a,β,9b,β))-dodecahydro-3a,6,6,9a-tetramethyl-naphthto(2,1-b)furan;
- (k) 0.05-10.0% amyl cinnamic aldehyde;
- (l) 0.05-0.9% benzyl propionate;
- (m) 0.05-0.30% L-1-methyl-4-isopropenyl-6-cyclohexene-2-one;
- 15 (n) 0.05-0.30% cis-3-hexenyl salicylate;
- (o) 0.05-0.70% 3,7-dimethyl-2,6-octadienal;
- (p) 0.05-2.00% 3-7-dimethyl-6-octen-1-ol;
- (q) 0.05-0.9% 2-methyl-3-(para-isopropylphenyl) propionaldehyde;
- (r) 0.05-0.9% 1-(2,6,6-trimethyl-2-cyclohexen-1-yl-2-buten-1-one);
- 20 (s) 0.005-0.05% decyl aldehyde Natural;
- (t) 0.05-0.3% 3-ethoxy-4-hydrobenzaldehyde;
- (u) 0.05-1.00% 2-methoxy-4-allyl phenol;

- (v) 0.05-0.30% 4-ethyl-alpha, alpha-dimethyl-benzenepropanal;
- (w) 0.05-6.0% 2-trans-3,7-dimethyl-2,6-octadien-8-ol;
- (x) 0.05-4.0% 3-(1-methyl-ethyl) bicyclo (2,2,1) hept-5-ene-2-carboxylic acid ethyl ester;
- 5 (y) 0.05-0.30% isobutyl quinoleine;
- (z) 0.05-2.0% 5-methyl-2-isopropyl cyclohexanol;
- (aa) 0.05-0.3% 4-isopropyl-1-methyl cyclohexan-3-one;
- (bb) 0.05-2.0% methyl ionone (mixture of isomers);
- (cc) 0.05-2.0% 3 methyl-4-(2,6,6-trimethyl-2-cyclohex-1-yl-3-buten-2-one;
- 10 (dd) 0.05-0.90% para-hydroxy phenyl butanone crystals;
- (ee) 0.05-4.0% 4-methyl-2-phenyl-3,6-dihydropyran;
- (ff) 0.05-0.30% 2 -(1,1-dimethylethyl)-4-methyl- cyclohexanol;
- (gg) 0.05-6.0% 1-methyl-4-isopropyl-1-cyclohexen-8-ol;
- (hh) 0.05-0.3% 4-hydroxy-3-methoxy benzaldehyde; and
- 15 (ii) 0.05-0.3% 2,4-diethoxy-5-methyl-pyrimidine;
- based on the total amount of fragrance component in a final cosmetic composition.

3. A cosmetic composition made with a fragrance enhancing component according to Claim 1.

20

4. A cosmetic composition made with a fragrance enhancing component according to Claim 2.

5. A cosmetic composition according to Claim 3 wherein the total amount

25 of the fragrance enhancing component added to the cosmetic composition is in the



range of 1.8-100% by weight based on the total fragrance component in the cosmetic composition.

6. A cosmetic composition according to Claim 4 wherein the total amount  
5 of the fragrance enhancing component added to the cosmetic composition is in the range of 1.8-100% by weight based on the total fragrance component in the cosmetic composition.

7. A cosmetic composition according to Claim 3 wherein the total amount  
10 of the fragrance enhancing component added to the cosmetic composition is in the range of 1.8-60% by weight based on the total fragrance component in the cosmetic composition.

8. A cosmetic composition according to Claim 4 wherein the total amount  
15 of the fragrance enhancing component added to the cosmetic composition is in the range of 1.8-60% by weight based on the total fragrance component in the cosmetic composition.

9. A cosmetic composition according to Claim 3 wherein the total amount  
20 of the fragrance enhancing component added to the cosmetic composition is in the range of 1.8-35% by weight based on the total fragrance component in the cosmetic composition.

25

10. A cosmetic composition according to Claim 4 wherein the total amount of the fragrance enhancing component added to the cosmetic composition is in the range of 1.8-35% by weight based on the total fragrance component in the cosmetic composition.

5

11. A cosmetic composition which provides a deodorant effect wherein the deodorant is made with a fragrance enhancing component itself made by combining at least three members selected from the group consisting of:

- (a) Cinnamon Leaf Oil Ceylan;
- 10 (b) Vetiver Oil Java;
- (c) Orange Oil Morocco;
- (d) Patchouli Oil;
- (e) Iso Methyl Cedryl Ketone A;
- (f) Pelargonyl;
- 15 (g) Cassis 345 B;
- (h) nonyl aldehyde;
- (i) decyl aldehyde;
- (j) 3aR-(3a,α,5a,α,9a,β,9b,β))-dodecahydro-3a,6,6,9a-tetramethyl-naphthto(2,1-b)furan;
- 20 (k) amyl cinnamic aldehyde;
- (l) benzyl propionate;
- (m) L-1-methyl-4-isopropenyl-6-cyclohexene-2-one;
- (n) cis-3-hexenyl salicylate;

- (o) 3,7-dimethyl-2,6-octadienal;
- (p) 3-7-dimethyl-6-octen-1-ol;
- (q) 2-methyl-3-(para-isopropylphenyl) propionaldehyde;
- (r) 1-(2,6,6-trimethyl-2-cyclohexen-1-yl)-2-buten-1-one;
- 5 (s) decyl aldehyde natural;
- (t) 3-ethoxy-4-hydrobenzaldehyde;
- (u) 2-methoxy-4-allyl phenol;
- (v) benzenepropanal, 4-ethyl-alpha, alpha-dimethyl;
- (w) 2-trans-3,7-dimethyl-2,6-octadien-8-ol;
- 10 (x) 3-(1-methyl-ethyl) bicyclo (2,2,1) hept-5-ene-2-carboxylic acid ethyl ester;
- (y) isobutyl quinoleine;
- (z) 5-methyl-2-isopropyl cyclohexanol;
- (aa) 4-isopropyl-1-methyl cyclohexan-3-one;
- (bb) methyl ionone (mixture of isomers);
- 15 (cc) 3-buten-2-one, 3 methyl-4-(2,6,6-trimethyl-2-cyclohex-1-yl);
- (dd) para-hydroxy phenyl butanone crystals;
- (ee) 4-methyl-2-phenyl-3,6-dihydropyran;
- (ff) cyclohexanol, 2 -(1,1-dimethylethyl)-4-methyl;
- (gg) 1-methyl-4-isopropyl-1-cyclohexen-8-ol};
- 20 (hh) 4-hydroxy-3-methoxy benzaldehyde; and
- (ii) pyrimidine, 2,4-diethoxy-5-methyl.

12. A cosmetic composition which provides a deodorant effect wherein the deodorant is made with a fragrance enhancing component itself made by combining at least three members selected from the group consisting of:
- (a) 0.5-1.0% Cinnamon Leaf Oil Ceylan;
  - 5 (b) 0.01-0.3% Vetiver Oil Java;
  - (c) 0.5-6.0% Orange Oil Morocco;
  - (d) 0.05-6.0% Patchouli Oil;
  - (e) 0.05-4.0% Iso Methyl Cedryl Ketone A;
  - (f) 0.05-6.0% Pelargonyl;
  - 10 (g) 0.05-0.9% Cassis 345 B;
  - (h) 0.005-0.05% Nonyl Aldehyde;
  - (i) 0.005-0.05% Decyl Aldehyde;
  - (j) 0.05-0.30% 3aR-(3a,α,5a,α,9a,β,9b,β))-dodecahydro-3a,6,6,9a-tetramethyl-naphthto(2,1-b)furan;
  - 15 (k) 0.05-10.0% amyl cinnamic aldehyde;
  - (l) 0.05-0.9% benzyl propionate;
  - (m) 0.05-0.30% L-1-methyl-4-isopropenyl-6-cyclohexene-2-one;
  - (n) 0.05-0.30% cis-3-hexenyl salicylate;
  - (o) 0.05-0.70% 3,7-dimethyl-2,6-octadienal;
  - 20 (p) 0.05-2.00% 3-7-dimethyl-6-octen-1-ol;
  - (q) 0.05-0.9% 2-methyl-3-(para-isopropylphenyl) propionaldehyde;
  - (r) 0.05-0.9% 2-buten-1-one, 1-(2,6,6-trimethyl-2-cyclohexen-1-yl);

- (s) 0.005-0.05% Decyl Aldehyde Natural;
  - (t) 0.05-0.3% 3-ethoxy-4-hydrobenzaldehyde;
  - (u) 0.05-1.00% 2-methoxy-4-allyl phenol;
  - (v) 0.05-0.30% benzenepropanal, 4-ethyl-alpha, alpha-dimethyl;
  - 5 (w) 0.05-6.0% 2-trans-3,7-dimethyl-2,6-octadien-8-ol;
  - (x) 0.05-4.0% 3-(1-methyl-ethyl) bicyclo (2,2,1) hept-5-ene-2-carboxylic acid ethyl ester;
  - (y) 0.05-0.30% isobutyl quinoleine;
  - (z) 0.05-2.0% 5-methyl-2-isopropyl cyclohexanol;
  - 10 (aa) 0.05-0.3% 4-isopropyl-1-methyl cyclohexan-3-one;
  - (bb) 0.05-2.0% methyl ionone (mixture of isomers);
  - (cc) 0.05-2.0% 3-buten-2-one, 3 methyl-4-(2,6,6-trimethyl-2-cyclohex-1-yl);
  - (dd) 0.05-0.90% para-hydroxy phenyl butanone crystals;
  - (ee) 0.05-4.0% 4-methyl-2-phenyl-3,6-dihydropyran;
  - 15 (ff) 0.05-0.30% 2-(1,1-dimethylethyl)-4-methyl-cyclohexanol;
  - (gg) 0.05-6.0% 1-methyl-4-isopropyl-1-cyclohexen-8-ol;
  - (hh) 0.05-0.3% 4-hydroxy-3-methoxy benzaldehyde; and
  - (ii) 0.05-0.3% 2,4-diethoxy-5-methyl pyrimidine;
- based on the amount of total fragrance component in the final cosmetic composition.

20

13. A cosmetic composition according to any one of Claims 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12, wherein the total fragrance component further comprises at least one member selected from the group consisting of:

(a) esters of salicylic acid selected from the group consisting of hexyl salicylate, hexenyl salicylate, isoamyl salicylate, benzyl salicylate and cyclohexyl salicylate;

(b) esters of cinnamic acid selected from the group consisting of amyl cinnamate, cinnamyl cinnamate and methyl cinnamate;

(c) esters selected from the group consisting of as allyl cyclohexane propionate, amyl benzoate, para-tertiarybutylcyclohexyl acetate, cedryl acetate, cedryl formate, dihydro-isojasmonate, ethylene brassylate, ethyl undecylenate, geranyl anthranilate, geranyl phenyl acetate, linalyl benzoate, benzyl acetate, linalyl acetate, vetiveryl acetate;

(d) aldehydes selected from the group consisting of amyl cinnamic aldehyde, cyclamen aldehyde, lillial, benzaldehyde, citronnellal, and hydroxycitronellal;

(e) alcohols selected from the group consisting of geraniol, linalool, nerol, phenyl ethyl alcohol, alpha terpineol, eugenol, isoeugenol, alpha-citronellol, dihydromyrcenol, aurantiol, cedrol, phenyl heptanol, phenol hexanol, alpha-santalol, undecavertol (4-methyl-3-decen-5-ol), and benzyl alcohol;

(f) ketones selected from the group consisting of benzophenone, dodecalactone, gamma-n-methyl ionone, delta-undecalactone, gamma-undecalactone, laevo-carvone, and beta-methylnaphthyl ketone; and

(g) nitromusks selected from the group consisting of musk ketone, musk tibetine, and musk indanone.

14. A cosmetic composition according to any one of Claims 3, 4, 5, 6, 7, 8, 9, 10, 11, or 12 further comprising an antiperspirant active added in an amount of 0.1-30% by weight based on the total weight of the composition.
- 5 15. A cosmetic composition according to any one of Claims 3, 4, 5, 6, 7, 8, 9, 10, 11, or 12 further comprising an antiperspirant active selected from the group comprising: aluminum chlorohydrate, aluminum chloride, aluminum sesquichlorohydrate, zirconyl hydroxychloride, aluminum-zirconium glycine complex (for example, aluminum zirconium trichlorohydrate gly, aluminum zirconium pentachlorohydrate gly, aluminum zirconium tetrachlorohydrate gly and aluminum zirconium octochlorohydrate gly), aluminum chlorohydrate PG, aluminum chlorohydrate PEG, aluminum dichlorohydrate PG, and aluminum dichlorohydrate PEG.
- 10 16. A cosmetic composition according to any one of Claims 3, 4, 5, 6, 7, 8, 9, 10, 11, or 12 further comprising a bacteriostat composition.
- 15 17. A cosmetic composition according to any one of 3, 4, 5, 6, 7, 8, 9, 10, 11, or 12 further comprising a bacteriostat composition selected from the group consisting of bacteriostatic quaternary ammonium compounds; 2-amino-2-methyl-1-propanol, cetyl pyridinium chloride; 2, 4, 4'-trichloro-2'-hydroxydiphenylether (Triclosan); N-(4-chlorophenyl)-N'-(3,4-dichlorophenyl)urea (Triclocarban); and bacteriostatic zinc salts.
- 20

18. A cosmetic composition according to Claim 3 made with a fragrance enhancing composition comprising 0.20-1.50% Cassis 345 B; 0.20-1.50% 2-buten-1-one, 1-(2,6,6-trimethyl-2-cyclohexen-1-yl); 0.20-1.50% para-hydroxy phenyl butanone crystals; 1.00-5.00% 3-(1-methyl-ethyl) bicyclo (2,2,1) hept-5-ene-2-carboxylic acid ethyl ester; 1.00-5.00% 4-methyl-2-phenyl-3,6-dihydropyrane based on the total weight of the fragrance component of the cosmetic composition, wherein the total amount of fragrance enhancing composition added is in the range of 2.60-14.50% based on the total weight of the fragrance component.
19. A cosmetic composition according to Claim 4 made with a fragrance enhancing composition comprising 0.40-.90% Cassis 345 B; 0.40-.90% 2-buten-1-one, 1-(2,6,6-trimethyl-2-cyclohexen-1-yl); 0.40-.90% para-hydroxy phenyl butanone crystals; 2.00-4.00% 3-(1-methyl-ethyl) bicyclo (2,2,1) hept-5-ene-2-carboxylic acid ethyl ester; 2.00-4.00% 4-methyl-2-phenyl-3,6-dihydropyrane based on the total weight of the fragrance component of the cosmetic composition, wherein the total amount of fragrance enhancing composition added is in the range of 5.2-10.7% based on the total weight of the fragrance component.
20. A cosmetic composition according to Claim 3 made with a fragrance enhancing composition comprising 0.02-0.50% 4-hydroxy-3-methoxy benzaldehyde; 0.02-0.50% pyrimidine, 2,4-diethoxy-5-methyl; 0.02-0.50% 3aR-(3a,α,5a,α,9a,β,9b,β))-dodecahydro-3a,6,6,9a-tetramethyl-naphthto(2,1-b)furan; 0.02-.5% benzenepropanal, 4-ethyl-α, α-dimethyl; 0.02-0.5% Vetiver Oil Java; 1.00-5.00% Iso Methyl Cedryl Ketone A; 1.00-5.00% 1-methyl-4-isopropyl-



1-cyclohexen-8-ol; 2.00-8.00 Patchouli Oil; and 4.00-12.00% amyl cinnamic aldehyde, based on the total weight of the fragrance component of the cosmetic composition, wherein the total amount of fragrance enhancing composition added is in the range of 8.10-32.50% based on the total weight of the fragrance component.

5

21. A cosmetic composition according to Claim 4 made with a fragrance enhancing composition comprising 0.05-0.30% 4-hydroxy-3-methoxy benzaldehyde; 0.05-0.30% pyrimidine, 2,4-diethoxy-5-methyl; 0.05-0.30% 3aR-(3a,α,5a,α,9a,β,9b,β))-dodecahydro-3a,6,6,9a-tetramethyl-naphthto(2,1-b)furan; 0.05-.30% benzenepropanal, 4-ethyl-α, α-dimethyl; 0.05-0.30% Vetiver Oil Java; 2.00-4.00% Iso Methyl Cedryl Ketone A; 2.00-4.00% 1-methyl-4-isopropyl-1-cyclohexen-8-ol; 3.00-6.00 Patchouli Oil; and 6.00-10.00% amyl cinnamic aldehyde, based on the total weight of the fragrance component of the cosmetic composition, wherein the total amount of fragrance enhancing composition added is in the range of 13.2-25.5% based on the total weight of the fragrance component.

15

22. A cosmetic composition according to Claim 3 made with a fragrance enhancing composition comprising 0.02-0.5% 2-buten-1-one, 1-(2,6,6-trimethyl-2-cyclohexen-1-yl); 0.02-0.5% cyclohexanol, 2-(1,1-dimethylethyl)-4-methyl-; 0.02-0.5 L-1-methyl-4-isopropenyl-6-cyclohexene-2-one; 0.20-1.20% benzyl propionate; 0.50-2.5% 2-methyl-3-(para-isopropylphenyl) propionaldehyde; 2.00-8.00% Orange Oil Morocco; 2.00-8.00% 3-7-dimethyl-6-octen-1-ol; 2.00-2.00-8.00% Pelargonyl, based on the total weight of the fragrance component of the cosmetic composition, wherein

20

the total amount of fragrance enhancing composition added is in the range of 6.76-29.20% based on the total weight of the fragrance component.

23. A cosmetic composition according to Claim 4 made with a fragrance enhancing composition comprising 0.05-0.30% 2-buten-1-one, 1-(2,6,6-trimethyl-2-cyclohexen-1-yl); 0.05-0.30% cyclohexanol, 2-(1,1-dimethylethyl)-4-methyl-; 0.05-0.30% L-1-methyl-4-isopropenyl-6-cyclohexene-2-one; 0.40-0.90% benzyl propionate; 1.00-2.00% 2-methyl-3-(para-isopropylphenyl) propionaldehyde; 3.00-6.00% Orange Oil Morocco; 3.00-6.00% 3-7-dimethyl-6-octen-1-ol; 3.00-6.00% Pelargonyl, based on the total weight of the fragrance component of the cosmetic composition, wherein the total amount of fragrance enhancing composition added is in the range of 10.6-21.8% based on the total weight of the fragrance component.

24. A cosmetic composition according to Claim 3 made with a fragrance enhancing composition comprising 0.002-0.1% nonyl aldehyde; 0.002-0.1% decyl aldehyde; 0.02-0.4% cis-3-hexenyl salicylate; 0.02-0.4% 3,7-dimethyl-2,6-octadienal; 0.90-1.20% 2-methyl-3-(para-isopropylphenyl) propionaldehyde; 0.8-2.40% 3-7-dimethyl-6-octen-1-ol; 0.8-2.40% methyl ionone (mixture of isomers), based on the total weight of the fragrance component of the cosmetic composition, wherein the total amount of fragrance enhancing composition added is in the range of 1.84-7.00% based on the total weight of the fragrance component.

25. A cosmetic composition according to Claim 4 made with a fragrance enhancing composition comprising 0.005-0.05% nonyl aldehyde; 0.005-0.05% decyl aldehyde; 0.05-0.30% cis-3-hexenyl salicylate; 0.05-0.30% 3,7-dimethyl-2,6-octadienal; 0.40-0.90% 2-methyl-3-(para-isopropylphenyl) propionaldehyde; 1.00-2.00% 3-7-dimethyl-6-octen-1-ol; 1.00-2.00% methyl ionone (mixture of isomers), based on the total weight of the fragrance component of the cosmetic composition, wherein the total amount of fragrance enhancing composition added is in the range of 2.5-5.6% based on the total weight of the fragrance component.

26. A cosmetic composition according to Claim 3 made with a fragrance enhancing composition comprising 0.02-0.4% isobutyl quinoleine; 0.02-0.40% 3aR-(3a,α,5a,α,9a,β,9b,β))-dodecahydro-3a,6,6,9a-tetramethyl-naphthto(2,1-b)furan; 0.4-1.20% Cinnamon Leaf Oil Ceylan; 0.4-1.20% 2-methoxy-4-allyl phenol; 1.00-5.00% Iso Methyl Cedryl Ketone A; and 2.00-7.00% Patchouli Oil, based on the total weight of the fragrance component of the cosmetic composition, wherein the total amount of fragrance enhancing composition added is in the range of 3.84-15.20% based on the total weight of the fragrance component.

27. A cosmetic composition according to Claim 4 made with a fragrance enhancing composition comprising 0.05-0.30% isobutyl quinoleine; 0.05-0.30% 3aR-(3a,α,5a,α,9a,β,9b,β))-dodecahydro-3a,6,6,9a-tetramethyl-naphthto(2,1-b)furan; 0.6-1.00% Cinnamon Leaf Oil Ceylan; 0.6-1.00% 2-methoxy-4-allyl phenol; 2.00-4.00% Iso Methyl Cedryl Ketone A; and 3.00-6.00% Patchouli Oil, based on the total weight of the fragrance component of the cosmetic composition, wherein the

total amount of fragrance enhancing composition added is in the range of 6.3-12.6% based on the total weight of the fragrance component.

28. A cosmetic composition according to Claim 3 made with a fragrance enhancing composition comprising 0.02-0.40% 4-isopropyl-1-methyl cyclohexan-3-one; 0.02-0.40% 3-ethoxy-4-hydrobenzaldehyde; 0.1-0.5% decyl aldehyde natural; 0.8-2.4% 5-methyl-2-isopropyl cyclohexanol; 0.8-2.4% 3-buten-2-one, 3 methyl-4-(2,6,6-trimethyl-2-cyclohex-1-yl; 2.00-7.00% 2-trans-3,7-dimethyl-2,6-octadien-8-ol, based on the total weight of the fragrance component of the cosmetic composition, wherein the total amount of fragrance enhancing composition added is in the range of 3.74-13.10% based on the total weight of the fragrance component.

29. A cosmetic composition according to Claim 4 made with a fragrance enhancing composition comprising 0.05-0.30% 4-isopropyl-1-methyl cyclohexan-3-one; 0.05-0.30% 3-ethoxy-4-hydrobenzaldehyde; 0.15-0.40% decyl aldehyde natural; 1.00-2.0% 5-methyl-2-isopropyl cyclohexanol; 1.00-2.00% 3-buten-2-one, 3 methyl-4-(2,6,6-trimethyl-2-cyclohex-1-yl; 3.00-6.00% 2-trans-3,7-dimethyl-2,6-octadien-8-ol, based on the total weight of the fragrance component of the cosmetic composition, wherein the total amount of fragrance enhancing composition added is in the range of 5.30-11.00% based on the total weight of the fragrance component.

30. A fragrance enhancing component according to Claim 1 made by combining five of the members.

31. A fragrance enhancing component according to Claim 2 made by combining five of the members.

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 99/04147

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 6 A61K7/32 A61K7/46

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 135 747 A (FARYNIARZ, J. R. ET AL.) 4 August 1992 see claims 1,3; example 2	1-4, 11-13,17
X,P	WO 98 35650 A (THE PROCTER AND GAMBLE CO.) 20 August 1998 see claim 1; tables 1-4	1-4, 11-13,17
X	US 5 114 717 A (KUZNITZ, M. ET AL.) 19 May 1992 see claim 1; table IV	1-4

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

7 June 1999

Date of mailing of the international search report

17/06/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Beyss, E

# INTERNATIONAL SEARCH REPORT

Information on patent family members

Intern. Application No

PCT/US 99/04147

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 5135747	A	04-08-1992	AT 121291 T	15-05-1995
			AU 652628 B	01-09-1994
			AU 1628292 A	19-11-1992
			CA 2068607 A,C	18-11-1992
			DE 69202107 D	24-05-1995
			DE 69202107 T	17-08-1995
			EP 0519531 A	23-12-1992
			ES 2072082 T	01-07-1995
			JP 2750044 B	13-05-1998
			JP 5148126 A	15-06-1993
			ZA 9203554 A	15-11-1993
WO 9835650	A	20-08-1998	US 5843881 A	01-12-1998
			AU 6159798 A	08-09-1998
US 5114717	A	19-05-1992	AT 108645 T	15-08-1994
			AU 652217 B	18-08-1994
			AU 1076792 A	13-08-1992
			CA 2060284 A,C	09-08-1992
			DE 69200242 D	25-08-1994
			DE 69200242 T	24-11-1994
			EP 0499304 A	19-08-1992
			ES 2056684 T	01-10-1994
			JP 2549963 B	30-10-1996
			JP 5058839 A	09-03-1993
			US 5198218 A	30-03-1993
			ZA 9200899 A	09-08-1993

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**